

MIC breakpoints ($\mu\text{g/ml}$):

Agent	Old (M100-S19)			Revised (M100-S20)		
	Susc	Int	Res	Susc	Int	Res
Cefazolin	≤ 8	16	≥ 32	≤ 1	2	≥ 4
Cefotaxime	≤ 8	16-32	≥ 64	≤ 1	2	≥ 4
Ceftizoxime	≤ 8	16-32	≥ 64	≤ 1	2	≥ 4
Ceftriaxone	≤ 8	16-32	≥ 64	≤ 1	2	≥ 4
Ceftazidime	≤ 8	16	≥ 32	≤ 4	8	≥ 16
Aztreonam	≤ 8	16	≥ 32	≤ 4	8	≥ 16

Disk diffusion breakpoints (mm):

Agent	Old (M100-S19)			Revised (M100-S20)		
	Susc	Int	Res	Susc	Int	Res
Cefazolin	≥ 18	15-17	≤ 14	NA	NA	NA
Cefotaxime	≥ 23	15-22	≤ 14	≥ 26	23-25	≤ 22
Ceftizoxime	≥ 20	15-19	≤ 14	≥ 25	22-24	≤ 21
Ceftriaxone	≥ 21	14-20	≤ 13	≥ 23	20-22	≤ 19
Ceftazidime	≥ 18	15-17	≤ 14	≥ 21	18-20	≤ 17
Aztreonam	≥ 22	16-21	≤ 15	≥ 21	18-20	≤ 17

NA = not available

CLSI M100-S20 (2010) recommended breakpoint changes for selected cephalosporins and aztreonam for *Enterobacteriaceae*.