

Appendiceal lesion cases, clues, and cautions

Amy Carpenter Aquino

May 2023—How to distinguish appendiceal diverticular disease and appendiceal polyps from mucinous neoplasms was just part of a CAP22 course on appendiceal lesions, led by Maryam Pezhouh, MD, MSc, of the University of California, San Diego, and Jacqueline Birkness-Gartman, MD, of Johns Hopkins University School of Medicine.

Their cases and comments covered mimics and more as they took attendees through inflammatory lesions, structural abnormalities, polyps, and gynecologic proliferations that involve the appendix. (Their comments on neoplasms and adenocarcinomas will be published separately.)

In her talk on inflammatory lesions of the appendix, Dr. Pezhouh, associate professor of pathology, reported on the case of a 45-year-old female who presented to the emergency room with abdominal pain, vomiting, and nausea. The patient's medical history included a renal transplant. Imaging was consistent with acute appendicitis and the patient underwent an appendectomy. **Fig. 1** reveals "the mucosa is basically ulcerated" with a lot of acute and chronic inflammation, Dr. Pezhouh said. In **Fig. 2** large cells with characteristic cytomegalovirus (CMV) nuclear inclusions also known as "owl eye" are seen. Immunostain (not shown) highlighted the infected cells.

"In acute appendicitis, we usually don't look for CMV," Dr. Pezhouh said, but it should be considered in patients with a history of solid organ transplant or HIV or who are very young or very old.

In the case of an eight-year-old boy who presented with abdominal pain and signs of acute appendicitis and who underwent an appendectomy, Dr. Pezhouh said inflammatory cells were seen in the lumen of the appendix and, on higher power, something else could also be seen in the lumen (**Fig. 3**). In **Fig. 4** is a cross-section of a worm in the lumen, and in **Fig. 5** is another worm or another section of the same worm, and eggs can be seen. It is *Enterobius vermicularis*, or pinworm, for which lateral alae are characteristic. Furthermore, typically *Enterobius vermicularis* eggs are plano-convex with one flattened side and one convex side. *Enterobius vermicularis* usually affects children ages seven to 11 and is an incidental finding, she said. Sometimes a mass of worms can cause obstruction in the GI tract or appendix, and occasionally pinworm can be associated with acute appendicitis. "Adult worms can also migrate to the lower genital tract and cause a granulomatous reaction, and the same goes for the appendix," Dr. Pezhouh said, though there were no granulomas in the case she reported.

In cases of a perforated appendix, sometimes clinicians treat the patient with antibiotics, discharge the patient, and perform an appendectomy later, as in the case of a 28-year-old male who presented with typical signs and symptoms of acute appendicitis and had surgery four weeks later, Dr. Pezhouh said. Sections of the appendix were inflamed, with a lot of lymphoid follicles and some focal acute inflammation, she said, but mostly chronic inflammation. Fairly well-formed granulomas are also seen in **Figs. 6** and **7**.

"When you see granulomas in the appendix, you know the differential is long," she said, one of which is interval (or delayed) appendicitis (when the patient has a ruptured appendix, is treated with antibiotics, and has an appendectomy four to eight weeks later). Its histologic features are variable and may include residual focal acute inflammation. Often seen is transmural chronic inflammation with lymphoid aggregates. Granulomatous inflammation is common (about 60 percent of the time). "You may see fibrosis often in the area that was perforated," post-healing, she said, and xanthogranulomatous appendicitis can be seen in a subset of cases.

Fig. 1




Fig. 2





Fig. 3



Granulomatous appendicitis is a rare diagnosis, she said—around 0.1 to two percent of appendectomies can have granulomas. “It usually resembles acute appendicitis” and usually doesn’t recur. “Only five to 10 percent of the patients can develop Crohn’s disease in the GI tract,” she said. It is usually idiopathic but can be caused by *Yersinia*, *Mycobacterium tuberculosis*, *Enterobius vermicularis*, a foreign body, interval appendicitis, and sarcoidosis, among other causes. “So remember the mimics and other etiologies to be considered before labeling the patient with Crohn’s disease whenever you see granulomas,” Dr. Pezhouh said.

Fig. 4




Fig. 5





Fig. 6



In **Fig. 8** is an example of Crohn's disease as seen in a longitudinal section of the appendix. It was the case of a 36-year-old female with a past medical history of Crohn's disease and refractory treatment who presented for ileocolonic resection, Dr. Pezhouh said. The inflammation is transmural. In **Fig. 9** granulomas are visible with focal areas of mild acute inflammation.

Gynecologic proliferations involving the appendix can take the form of endometriosis, endosalpingiosis, and decidual lesions, said Dr. Birkness-Gartman, assistant professor of pathology at Johns Hopkins. Endometriosis consists of endometrial-type glands and often has associated endometrial stroma, and there can be hemorrhage and hemosiderin-laden macrophages, she said. Endosalpingiosis consists of glands or cystic structures lined by tubal-type epithelium without associated stroma. "And decidual lesions can consist of decidualized endometriosis where you have decidualized stroma with some retained glands," she said, "or deciduosis where you do not have glands."

Fig. 7






Fig. 8.



Dr. Birkness-Gartman presented the case of a 40-year-old woman with known endometriosis who underwent a diagnostic laparoscopy, which revealed endometriosis involving both ovaries, the appendix, and the colonic serosa. The patient underwent a hysterectomy with bilateral salpingo-oophorectomy and appendectomy. In **Fig. 10** is a fibrotic-appearing lesion involving the appendix (arrows). In **Fig. 11** are glands that appear to be infiltrating the muscularis propria.

In **Fig. 12** is “the typical histology of endometriosis,” she said, noting the endometrial-type glands, “and one of the clues to the diagnosis is the spindled endometrial-type stroma surrounding the glands.”


Fig. 9**Fig. 10**

Case courtesy of Dr. Monica Butcher, Johns Hopkins Atlas of Appendiceal Pathology


Fig. 11

Case courtesy of Dr. Monica Butcher, Johns Hopkins Atlas of Appendiceal Pathology

Noor, et al., reported finding in a review of cases that appendiceal endometriosis affects women with a median age of 34, Dr. Birkness-Gartman said, “and this could present with acute appendicitis, intussusception, or GI bleeding” (Noor M, et al. *Hum Pathol.* 2019;92:101–106). They found that 73 percent of patients with appendiceal endometriosis had extra-appendiceal involvement, “but interestingly,” she said, “those who presented with acute appendicitis were less likely to have extra-appendiceal involvement” (22 percent). Histologically, appendiceal endometriosis typically involves the muscularis propria but can involve the serosa, she said. Glands can have an infiltrative-appearing growth pattern, “which could raise concern for adenocarcinoma.” However, endometrial-type stroma, hemorrhage, and hemosiderin-laden macrophages are clues to the correct diagnosis, she said. IHC can be helpful because endometrial-type glands will label for CK7 and ER, and the stroma will label for CD10 and ER.



Fig. 12

Case courtesy of Dr. Monica Butcher, Johns Hopkins Atlas of Appendiceal Pathology


Fig. 13

Case courtesy of Dr. Lysandra Voltaggio


Endometriosis with intestinal metaplasia is one pitfall, Dr. Birkness-Gartman said. “It can potentially mimic a low-grade appendiceal mucinous neoplasm or LAMN.” Associated endometrial-type stroma is a clue, she said. Endometrial-type glands with intestinal metaplasia will label for the intestinal marker. “Luckily, the endometrial stroma retains the expected labeling pattern and is positive for ER and CD10,” she said. In a high-magnification image of endometriosis with intestinal metaplasia (**Fig. 13**), the endometrial-type stroma surrounding the glands has a spindled appearance and “is a clue to the correct diagnosis in this case” (Vyas M, et al. *Pathol Res Pract.* 2017;213[1]:39-44).

Fig. 14**Fig. 15**

Dr. Birkness-Gartman presented the case of a 40-year-old woman with a history of ampullary adenocarcinoma who underwent a Whipple procedure with en bloc ileocecectomy. Incidental glandular-type structures are seen in **Fig. 14** in the periappendiceal adipose tissue and at higher magnification there is no associated endometrial-type stroma (**Fig. 15**). It is lined by cuboidal epithelial cells that have cilia on the luminal surface. Appendiceal endosalpingiosis is an incidental finding in this case, she said.

Fig. 16

Case courtesy of Dr. Lysandra Voltaggio, Johns Hopkins Atlas of Appendiceal Pathology

Fig. 17

Case courtesy of Dr. Lysandra Voltaggio, Johns Hopkins Atlas of Appendiceal Pathology

Noor, et al., reported finding that the median age of patients with appendiceal endosalpingiosis was 45, and 36 percent of the patients in their study presented with acute appendicitis. “Extra-appendiceal involvement can be

seen in 36 percent of patients, but it's typically not in patients presenting with acute appendicitis," Dr. Birkness-Gartman said. Histologically, appendiceal endosalpingiosis can involve the muscularis propria or serosa, typically in minute foci, she said. "And we see glands or cystic structures lined by tubal-type epithelium that lack the associated endometrial-type stroma and hemosiderin."

In another case, a 30-year-old woman who was 13 weeks pregnant presented with acute appendicitis and was incidentally found to have an appendiceal mass, Dr. Birkness-Gartman said. In **Fig. 16** a large eosinophilic proliferation can be seen extending from the serosal surface, "almost the whole way to the luminal surface of the appendix," she said. In an image taken on the serosal surface (**Fig. 17**), large polygonal cells with abundant pink cytoplasm can be seen with associated acute serositis (at top), she said. On IHC, the cells were positive for PR and CD10.

"So we signed this out as exuberant deciduosis" to indicate that it was forming a mass, Dr. Birkness-Gartman said, "and we noted that it was associated with acute appendicitis and serositis." They mentioned it was positive for PR and CD10 as expected for deciduosis. Additional immunostains showed the decidualized cells were negative for CD68, S100, and AE1/AE3, ruling out some of their differential diagnoses, she said.

Appendiceal deciduosis affects women (median age 31) who may be pregnant or taking oral contraceptives, Dr. Birkness-Gartman said. Acute appendicitis is the most common presentation (71 percent), and extra-appendiceal involvement is seen in 43 percent of cases. Histologically, the serosa is typically involved, "but this may spread transmurally and may even obliterate the appendix in some cases," she said. The cells are eosinophilic with voluminous cytoplasm and on IHC will label for PR and CD10 (Noor M, et al. *Hum Pathol.* 2019;92:101-106).

In the differential diagnosis is xanthogranulomatous appendicitis, but the cytoplasm has a foamy quality, "which helps you morphologically," and it will be positive for CD68. Also in the differential is granular cell tumor, which, while it too has voluminous pink cytoplasm, tends to have a more granular quality to the cytoplasm, she said, and it labels for CD68 and S100. Their differential includes metastatic carcinoma and mesothelioma, "both of which would label for keratins," and other neoplasms.

In Dr. Pezhouh's discussion of structural abnormalities of the appendix, she presented the case of a 50-year-old male with appendiceal irregularity, a mass lesion on imaging, and a medical history of diabetes and cystic fibrosis. He underwent an appendectomy. In **Fig. 18** the lumen is dilated and "filled with pink eosinophilic mucus," she said. In **Fig. 19** some of the crypts are enlarged and have inflammatory cells in them mixed with eosinophilic secretions.

This was a case of appendiceal mucocoele in the setting of a cystic fibrosis patient, Dr. Pezhouh said. Appendiceal mucocoele is a clinical term, not a histologic term, she noted, and describes a "grossly dilated appendix filled with mucus." Its causes are a hyperplastic polyp (similar to a colonic polyp); mucinous neoplasm, most commonly LAMN and high-grade appendiceal mucinous neoplasm; and simple mucocoele from endometriosis obstruction or inspissated eosinophilic mucus in cases of patients with cystic fibrosis.

Another case Dr. Pezhouh presented was that of a 45-year-old male patient with intermittent abdominal pain and a low-grade fever who had an appendectomy after imaging and clinical findings pointed to acute appendicitis. In **Fig. 20** there was focally acute appendicitis and the lumen of the appendix had lymphoid aggregates, but it was otherwise "devoid of acute appendicitis," she said, noting diverticular formation, on right.

Appendiceal diverticulosis is uncommon (around 0.004 to two percent of appendectomies), "but 22 percent of patients with cystic fibrosis will present with diverticulosis or diverticular disease because of the pressure in the lumen," Dr. Pezhouh said. Acquired appendiceal diverticulosis, in which there is no muscularis propria, is more common, she said, while in the congenital version of the disease, muscularis propria surrounds the area of diverticulosis. "It's basically due to increased intraluminal pressure and wall weakness in the GI tract," and is the same in the colon, she said. It can cause diverticulitis or inflammation and, if so, will resemble appendicitis. "It can also perforate, and if it perforates you can have high mortality. If it's associated with LAMN, it also, when perforated, can cause pseudomyxoma peritonei."

Complicated diverticulosis, when it's associated with mucus cysts and enlargement of the appendix, can resemble LAMN, Dr. Pezhouh said. Retention of the normal mucosal architecture with lamina propria is one indication of diverticular disease, she said. Other clues in favor of diverticulosis: crypt architecture is maintained, crypts are arranged in a regular array, and there is a lack of nuclear abnormalities. (See **Fig. 21.**)

Appendiceal polyps are similar morphologically to those that occur elsewhere in the colon, Dr. Birkness-Gartman said. "So you can see the typical tubular, tubulovillous, and villous adenomas, hyperplastic polyps, sessile serrated adenomas—which the WHO now calls sessile serrated lesions—and traditional serrated adenoma." They're treated with resection with negative margins, "as long as they have not yet developed adenocarcinoma because they can be precursor lesions."

Fig. 18






Fig. 19




In distinguishing appendiceal polyps from LAMN, she said, "you should see [in appendiceal polyps] preserved lamina propria and muscularis mucosae, and this should not show pushing invasion where you lose those structures and get underlying fibrosis." The cytologic features of a tubulovillous adenoma involving the appendix are seen in **Fig. 22**—"typical conventional low-grade dysplasia with elongation, crowding, and hyperchromasia of the nuclei," Dr. Birkness-Gartman noted, as well as "retained architecture of the lamina propria and muscularis mucosae, which is why I would not diagnose this as a LAMN."

Fig. 20




She presented the case of a 45-year-old woman with an elongated, polypoid mass at the appendiceal orifice who was treated with an ileocecectomy. The bisected lesion revealed a core of adipose tissue, and in **Fig. 23** the villiform structure and pink epithelium is seen. On the right in Fig. 23 "we see the diagnostic feature, which is ectopic crypt foci with little crypts sort of budding off of the villi, and they're not oriented with their bases toward the muscularis mucosae."

Fig. 21




In this case, there is a traditional serrated adenoma and, secondly, an inverted appendix, Dr. Birkness-Gartman said. “We had a neoplasm that arose in the appendix and it essentially served as a lead point, and the appendix turned inside out. It became inverted.” Appendiceal inversion has been associated with various neoplasms and with endometriosis—“basically anything that can serve as a lead point for intussusception,” she said (Birkness J, et al. *Histopathology*. 2019;74[6]:853-860).

Fig. 22. Tubulovillous adenoma



Dr. Birkness-Gartman presented the case of a 60-year-old woman with acute appendicitis, whose slides she and colleagues received in consultation about whether it could be a LAMN. In **Fig. 24** the epithelium looks “frisky,” she noted. The area indicated with an asterisk on the left side turned out to be acute appendicitis with abscess formation at higher magnification, she said. “So whatever this lesion is, it’s arising in the background of appendicitis.” In **Fig. 25** the lamina propria and muscularis mucosae are seen. “So I think this is not a LAMN based upon that architecture,” Dr. Birkness-Gartman said, noting also the hyperplastic or serrated-looking architecture, “which was more pronounced near the luminal surface.”

Fig. 23




Photos courtesy of Dr. Lysandra Voltaggio

Fig. 24



Fig. 23




Photos courtesy of Dr. Lysandra Voltaggio

Fig. 24



It is an example of post-inflammatory mucosal hyperplasia, a benign finding associated with prior appendicitis and one that can also be seen in cases of interval appendicitis, she said (Hissong E, et al. *Mod Pathol.* 2020;33[5]:953–961). “It lacks the architectural features of LAMN; specifically, it does not show a pushing border.”□

Fig. 25



Amy Carpenter Aquino is CAP TODAY senior editor. Images are courtesy of Drs. Pezhouh and Birkness-Gartman, as well as Monica Butcher, MD, and Lysandra Voltaggio, MD, whose images published in the preceding article appear in the Johns Hopkins Pathology Atlas iPad app on appendiceal pathology, released in 2022 (<https://apps.pathology.jhu.edu/all>). Dr. Birkness-Gartman is first author on the app; Toby Cornish, MD, PhD,

Norman Barker, MA, MS, and Ralph Hruban, MD, are series editors.